



**Institute of Biodiversity and Environmental Conservation**

**Pollination Guilds of Aroids in Mulu National Park**

**Chai Shong Kian**

**Master of Science  
2020**

# Pollination Guilds of Aroids in Mulu National Park

Chai Shong Kian

A thesis submitted

In fulfilment of the requirements for the degree of Master of Science

(Plant Science)

Institute of Biodiversity and Environmental Conservation

UNIVERSITI MALAYSIA SARAWAK

2020

## DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Malaysia Sarawak. Except where due acknowledgement has been made, the work is that of the author alone. The thesis has not been accepted for any degree and is not concurrently submitted in candidature of any other degree.

.....

Signature

Name: Chai Shong Kian

Matric No.: 17020182

Institute of Biodiversity and Environmental Conservation

Universiti Malaysia Sarawak

Date:

## **ACKNOWLEDGEMENT**

Firstly, I would like to express sincere gratitude to my supervisor, Assoc. Prof. Dr Wong Sin Yeng who encouraged me to continue M.Sc. Without her continuous guidance throughout all the stages of the project, it would be impossible to complete the project. Her patience, motivation and knowledge helped me to achieve the best possible for the project.

My sincere thanks also go to the staff of Borsarmulu Park Management and Sarawak Forestry Corporation in Mulu. All of them were very friendly and glad to assist me during my stay. Mr Hein Gerstner, the Park Manager, treated me like one of his staff, always concerned about me and granted me special right to have cheaper Wi-Fi rate, and free access to all tours available in the park.

Next, I would like to acknowledge the Ministry of Education of Malaysia for funding this project through Vote No. NRGs/1089/2013-(03). Fieldwork associated with this project was most recently under Sarawak Forestry Department Permission to Conduct Research on Biological Resources Permit No. (123)JHS/NCCD/600-7/2/107 and Park Permit No. WL64/2018.

Last but not least, I would like to thank my family members especially my mother who always supports me and listens to my heartfelt mumblings. Her advice makes me more cautious to work safely in the field.

## ABSTRACT

The aroid flora of Borneo is one of the richest on the planet, comprising at least 800 species with many highly localized endemics. Not surprisingly, most of our knowledge of these taxa is limited to naming taxonomic novelties but beyond this, our knowledge on the pollination biology of Bornean aroids remains poor. Previous pollination research on aroid taxa on Borneo, revealed two pollination guilds: Diptera (Drosophilidae) and Coleoptera (Chrysomelidae, Hydrophilidae, and Staphylinidae) which are by far the major pollinating groups. This study was conducted at Gunung Mulu National Park, an area which hold one of the richest aroid diversity on Borneo, and aimed to determine pollinators of 21 selected aroids belonging to Tribe Aglaonemateae, Anadendrieae, Colocasieae, Homalomeneae, Monstereae, Potheae, Schismatoglottideae, and Thomsonieae. Extensive field observations on floral biology including stages of floral development, floral traits and rewards, identification of insect pollinators, and their behaviour(s) were carried out. The total anthetic time of most unisexually flowered taxa (*Alocasia*, *Aglaonema*, *Homalomena*, and Schismatoglottideae) lasted for two days, except *Amorphophallus* which ranged two to five days. In bisexually flowered taxa, *Scindapsus* completed anthesis in two days but anthesis lasted four days in *Anadendrum* and longer in *Lasia* and *Pothos*, which ranged 7-18 days. All investigated aroid taxa emitted floral odour during pistillate anthesis with several taxa (*Anadendrum*, *Bucephalandra*, *Homalomena*, *Pothos*, and *Schismatoglottis pellucida*) extended odour production until the end of anthesis. Some taxa (*Amorphophallus* and *Pothos*) do not restrict access to the pistillate flowers during pistillate anthesis. Diptera flies are the most possible pollinators for *Alocasia*, *Aglaonema*, *Homalomena*, *Pothos*, Schismatoglottideae and *Scindapsus*; Coleoptera beetles are the pollinators for *Amorphophallus*, *Anadendrum*, *Homalomena*, *Lasia*, and *Schismatoglottis*; Thysanoptera

(thrips) are the pollinators of *Anadendrum*; Hemiptera are pollinators of *Lasia* and *Pothos*; Hymenoptera are pollinators of *Alocasia sarawakensis* and *Amorphophallus hewittii*. Floral rewards for pollinators included stigmatic secretion, floral tissues, and provision of mating sites.

**Keywords:** *Amorphophallus*, *Anadendrum*, anthesis, *Lasia*, pollinators

## ***Kumpulan Pendebungaan Keladi Hutan di Taman Negara Mulu***

### **ABSTRAK**

*Flora keladi hutan Borneo adalah salah satu yang terkaya di planet ini, dengan sekurang-kurangnya 800 spesies yang terdiri daripada banyak endemik tempatan. Tidak menghairankan, kebanyakan pengetahuan tentang keladi hutan tersebut adalah terhadap kepada penamaan taksonomi baru tetapi luar daripada ini, pengetahuan tentang biologi pendebungaan keladi hutan dari Borneo masih kurang. Penyelidikan pendebungaan tentang keladi hutan di Borneo sebelum ini, mendedahkan dua kumpulan pendebunga: Diptera (*Drosophilidae*) dan Coleoptera (*Chrysomelidae*, *Hydrophilidae*, dan *Staphylinidae*) yang merupakan kumpulan pendebunga utama. Kajian ini dijalankan di Taman Negara Gunung Mulu, salah satu kawasan yang mempunyai kepelbagaian keladi hutan terkaya di Borneo, dan bertujuan untuk menentukan pendebunga untuk 21 keladi hutan terpilih daripada Tribe *Aglaonemateae*, *Anadendrieae*, *Colocasieae*, *Homalomeneae*, *Monstereae*, *Potheae*, *Schismatoglottideae*, dan *Thomsonieae*. Pemerhatian lapangan secara meluas mengenai biologi bunga termasuk peringkat perkembangan bunga, sifat bunga dan ganjaran, pengenalpastian pendebunga serangga, dan tingkah laku mereka telah dijalankan. Jumlah masa antesis untuk kebanyakan keladi hutan berbunga uniseks (*Alocasia*, *Aglaonema*, *Homalomena*, dan *Schismatoglottideae*) berlangsung selama dua hari, kecuali *Amorphophallus* yang berjulat antara dua hingga lima hari. Untuk keladi hutan berbunga biseks, *Scindapsus* menyempurnakan proses antesis dalam dua hari tetapi antesis bertahan selama empat hari di *Anadendrum* dan lebih lama di *Lasia* dan *Pothos*, yang berjulat antara 7-18 hari. Semua taxa keladi hutan mengeluarkan bau bunga semasa antesis pistillate dengan beberapa keladi hutan (*Anadendrum*, *Bucephalandra*, *Homalomena*, *Pothos*, dan*

*Schismatoglottis pellucida*) mengeluarkan bau bunga sehingga antesis berakhir. Sesetengah keladi hutan (*Amorphophallus* dan *Pothos*) tidak menghadkan akses ke bunga jantina semasa antesis pistillate. Lalat Diptera berkemungkinan besar adalah pendebunga untuk *Alocasia*, *Aglaonema*, *Homalomena*, *Pothos*, *Schismatoglottideae* dan *Scindapsus*; Kumbang Coleoptera adalah pendebunga untuk *Amorphophallus*, *Anadendrum*, *Homalomena*, *Lasia*, dan *Schismatoglottis*; Thysanoptera (thrips) adalah pendebunga untuk *Anadendrum*; Hemiptera adalah pendebunga untuk *Lasia* dan *Pothos*; Hymenoptera adalah pendebunga untuk *Alocasia sarawakensis* dan *Amorphophallus hewittii*. Ganjaran bunga untuk pendebunga termasuk cecair stigma, tisu bunga, dan penyediaan tapak mengawan.

**Kata kunci:** *Amorphophallus*, *Anadendrum*, antesis, *Lasia*, pendebunga



## TABLE OF CONTENTS

	Page
<b>DECLARATION .....</b>	<b>i</b>
<b>ACKNOWLEDGEMENT .....</b>	<b>ii</b>
<b>ABSTRACT .....</b>	<b>iii</b>
<b><i>ABSTRAK</i> .....</b>	<b>v</b>
<b>TABLE OF CONTENTS .....</b>	<b>vii</b>
<b>LIST OF TABLES.....</b>	<b>xiii</b>
<b>LIST OF FIGURES.....</b>	<b>xv</b>
<b>LIST OF ABBREVIATIONS.....</b>	<b>xxviii</b>
<b>CHAPTER 1: INTRODUCTION.....</b>	<b>1</b>
1.1 Background.....	1
1.2 Problem statement.....	3
1.3 Objectives .....	3
<b>CHAPTER 2: LITERATURE REVIEW.....</b>	<b>4</b>
2.1 Araceae .....	4
2.1.1 Life forms .....	4
2.1.2 Vegetative morphology .....	7
2.1.3 Floral morphology .....	7
2.1.4 Fruits and seeds .....	11

2.1.5	Seed dispersal .....	13
2.2	Araceae on Borneo.....	15
2.2.1	Subfamily Pothoideae.....	16
2.2.2	Subfamily Monsteroideae.....	17
2.2.3	Subfamily Lasioideae .....	18
2.2.4	Subfamily Aroideae.....	19
2.3	Pollination study of Araceae.....	24
2.3.1	Subfamily Pothoideae.....	27
2.3.2	Subfamily Monsteroideae.....	27
2.3.3	Subfamily Lasioideae .....	28
2.3.4	Subfamily Aroideae.....	29
<b>CHAPTER 3:</b>	<b>MATERIALS AND METHODS .....</b>	<b>35</b>
3.1	Study area .....	35
3.2	Plant species.....	38
3.3	Survey of flowering populations.....	41
3.4	Flowering pattern .....	44
3.5	Floral developmental stages.....	44
3.6	Insect visitors and pollinators .....	47
3.7	Reproductive system.....	48
<b>CHAPTER 4:</b>	<b>RESULTS.....</b>	<b>49</b>
4.1	<i>Aglaonema</i> .....	49

4.1.1	Flowering pattern.....	49
4.1.2	Timing of anthesis and floral changes .....	54
4.1.3	Floral traits.....	57
4.1.4	Floral rewards .....	57
4.1.5	Pollinators and their behaviours .....	58
4.1.6	Reproductive system .....	63
4.2	<i>Alocasia</i> .....	64
4.2.1	Flowering pattern.....	64
4.2.2	Timing of anthesis and floral changes .....	67
4.2.3	Floral traits.....	67
4.2.4	Floral rewards .....	68
4.2.5	Pollinators and their behaviours .....	68
4.2.6	Reproductive system .....	69
4.3	<i>Amorphophallus</i> .....	69
4.3.1	Flowering pattern.....	69
4.3.2	Timing of anthesis and floral changes .....	75
4.3.3	Floral traits.....	76
4.3.4	Floral rewards .....	77
4.3.5	Pollinators and their behaviours .....	77
4.3.6	Reproductive system .....	79
4.4	<i>Anadendrum</i> .....	80

4.4.1	Flowering pattern.....	80
4.4.2	Timing of anthesis and floral changes .....	83
4.4.3	Floral traits.....	83
4.4.4	Floral rewards .....	83
4.4.5	Pollinators and their behaviours .....	84
4.4.6	Reproduction system .....	84
4.5	<i>Bucephalandra</i> .....	85
4.5.1	Flowering pattern.....	85
4.5.2	Timing of anthesis and floral changes .....	87
4.5.3	Floral traits.....	87
4.5.4	Floral rewards .....	88
4.5.5	Pollinators and their behaviours .....	88
4.5.6	Reproductive system .....	88
4.6	<i>Homalomena</i> .....	89
4.6.1	Flowering pattern.....	89
4.6.2	Timing of anthesis and floral changes .....	90
4.6.3	Floral traits.....	99
4.6.4	Floral rewards .....	99
4.6.5	Pollinators and their behaviours .....	100
4.6.6	Reproductive system .....	102
4.7	<i>Lasia</i> .....	102

4.7.1	Flowering pattern.....	102
4.7.2	Timing of anthesis and floral changes .....	103
4.7.3	Floral traits.....	107
4.7.4	Floral rewards .....	107
4.7.5	Pollinators and their behaviours .....	107
4.7.6	Reproduction system .....	109
4.8	<i>Pothos</i> .....	110
4.8.1	Flowering pattern.....	110
4.8.2	Timing of anthesis and floral changes .....	110
4.8.3	Floral traits.....	116
4.8.4	Floral rewards .....	116
4.8.5	Pollinators and their behaviours .....	116
4.8.6	Reproduction system .....	118
4.9	<i>Schismatoglottis</i> .....	118
4.9.1	Flowering pattern.....	118
4.9.2	Timing of anthesis and floral changes .....	120
4.9.3	Floral traits.....	135
4.9.4	Floral rewards .....	135
4.9.5	Pollinators and their behaviours .....	136
4.9.6	Reproductive system .....	139
4.10	<i>Scindapsus</i> .....	139

4.10.1	Flowering pattern.....	139
4.10.2	Timing of anthesis and floral changes .....	139
4.10.3	Floral traits.....	142
4.10.4	Floral rewards .....	142
4.10.5	Pollinators and their behaviours .....	142
4.10.6	Reproductive system .....	143
<b>CHAPTER 5:</b>	<b>DISCUSSION .....</b>	<b>144</b>
5.1	Flowering duration.....	144
5.2	Floral traits .....	146
5.3	Floral rewards .....	148
5.4	Pollination guilds .....	149
5.5	Reproductive system.....	154
<b>CHAPTER 6:</b>	<b>CONCLUSION AND RECOMMENDATIONS .....</b>	<b>157</b>
6.1	Conclusion .....	157
6.2	Recommendations.....	160
<b>REFERENCES</b>	<b>.....</b>	<b>161</b>
<b>APPENDICES</b>	<b>.....</b>	<b>182</b>

## LIST OF TABLES

	<b>Page</b>
Table 3.1      A checklist of aroid species in Mulu National Park. 1 non native. * taxa included in this study. 95 taxa including one non-native.	39
Table 3.2      Life form/ecology, habitat, and location of the 21 investigated aroid species.	42
Table 3.3      Number of replicates (inflorescences utilised) for field observation (timing of anthesis, floral traits, rewards, insect visitors and their behaviour), insects captured and pollination treatments (autonomous self- and open- pollination).	43
Table 3.4      Measurement of spathe (widest point during pistillate anthesis) and spadix for the 21 studied aroid species.	45
Table 4.1      The timing of anthetic stages (sunrise time: 0600-0630 h, sunset time: 1830-1900h), floral odour, restricted access to flower (during pistillate anthesis), duration of fruit maturity and dispersal agent of the investigated taxa. (n = no. of replicates)	55
Table 4.2      Mean $\pm$ SE, range and percentage of occurrence of insect families captured for each species during anthesis. (n=no. of replicates)	59

Table 4.3	Mean $\pm$ SE of pistillate flowers per inflorescences and fruits per infructescence, and fruit set percentage of the aroid species treated with open pollination treatment. (n=no. of replicates)	63
-----------	--	----



## LIST OF FIGURES

	<b>Page</b>
Figure 3.1	36
<p>Geological map of Mulu National Park, the area of study is part of the Melinau limestone formation. Adapted from A guide to the karst geomorphology of Mulu, by A. R. Farrant, 2009, Retrieved from <a href="http://www.mulucaves.org/wordpress/science/the-geomorphology-of-mulu">http://www.mulucaves.org/wordpress/science/the-geomorphology-of-mulu</a>.</p>	
Figure 3.2	36
<p>Map of trails near HQ of Mulu National Park, the trails of this study included Deer cave walk, Botany Walk, Canopy walk, Paku waterfall and the night walk. Retrieved and edited from <a href="http://www.etawau.com/Geography/Sarawak/04_Miri/Miri/MuluTour.htm">http://www.etawau.com/Geography/Sarawak/04_Miri/Miri/MuluTour.htm</a>.</p>	
Figure 3.3	37
<p>Map of the forest type of Mulu National park, the area of study included area from HQ to Deer Cave and Camp 5, the forest type of the study area are riverine forest and lowland limestone forest. Adapted from A Guide to Gunung Mulu National Park, H. P. Hazebroek &amp; A. K. Abang Morshidi, 2002, Borneo: N.H.P.B.</p>	
Figure 4.1	50
<p><i>Aglaonema nitidum</i> (A) Whole plant with inflorescences and infructescences. (B) Post anthetic inflorescence, spathe closed back (C) Immature fruits. (D) About to ripe fruits in pink and ripening fruits in red.</p>	

- Figure 4.2    Inflorescence of *Aglaonema nitidum* at different stages. (A-C)    51  
Pistillate anthesis, spathe gap widely opened, flies entered. (D)  
Inflorescence before staminate anthesis, spathe gap closed back.  
(E) Staminate anthesis. (F) Receptive pistils during pistillate  
anthesis, *Colocasiomyia* flies (Drosophilidae) at sterile interstice.  
(G) *Colocasiomyia* flies (Drosophilidae) collecting pollen grains.  
(H) *Colocasiomyia* flies (Drosophilidae) moved onto pistils,  
unknown flies at sterile interstice during pistillate anthesis.
- Figure 4.3    *Aglaonema simplex* (A) Adult plants with inflorescences and    52  
infructescences. (B) Immature fruits in yellow and ripened fruits in  
red. (C) Inflorescence during pre-anthesis and post anthesis (spathe  
and staminate zone abscised).
- Figure 4.4    Inflorescence of *Aglaonema simplex* at different stages. (A-C)    53  
Pistillate anthesis, spathe gap widely opened. (B) Unknown flies  
arrived before *Colocasiomyia* flies (Drosophilidae) arrive. (C-E)  
*Colocasiomyia* flies arrived and stayed throughout anthesis,  
pistillate flowers turned deep yellow post pistillate anthesis. (E)  
Staminate anthesis, spathe gap closed back. (F) *Colocasiomyia* flies  
move onto pistillate flowers. (G) Unknown flies mate at staminate  
zone. (H) *Colocasiomyia* flies collect pollen. (I) *Cycreon* beetles  
(Hydrophilidae) at pistillate anthesis.
- Figure 4.5    *Alocasia sarawakensis* (A) Multiple inflorescences and    65  
infructescences per plant. (B) Dead stingless bees trapped at

pistillate zone. (C) Weaver ant (*Oecophylla* sp.) preying on a *Colocasiomyia* fly. (D-E) Bulbuls (D) and Sunbird (E) consuming ripened fruits.

- Figure 4.6    Inflorescence of *Alocasia sarawakensis* at various stages. (A)    66  
Pistillate anthesis. (B) Pre staminate anthesis, spathe curve backward. (C) Flies (Drosophilidae) move and mate on pistillate flowers and lower inner side of the spathe. (D) Flies at pistillate zone (Top view). (E) Stingless bees and unknown flies collecting pollen on the staminate zone. (F) Flies (Drosophilidae) collecting pollen on the staminate zone (Top view).
- Figure 4.7    *Amorphophallus hewittii* (A) Vegetative form, distinguished by    70  
pattern of soldier camouflage cloth on the petiole. (B) Floral zone (top staminate, bottom pistillate). (C) String form pollen released. (D) Ripened fruits, white colour before ripe.
- Figure 4.8    Inflorescence of *Amorphophallus hewittii* at various stages. (A-D)    71  
Changes of inflorescence throughout anthesis (front view), (A-C) spathe expanding (pistillate anthesis), (D) spathe collapsing (Staminate anthesis). (E-H) Changes of inflorescence throughout anthesis (top view), (E-G) Spathe expanding (pistillate anthesis), (H) Spathe collapsing. (I-L) Zoom view of spadix throughout anthesis, (I-K) Liquid condensed from appendix, started from bottom to tip, floral odour more intense as more liquid condensed

(Pistillate anthesis). (L) Appendix dried, smell weakened  
(Staminate anthesis).

- Figure 4.9 Insects on different parts of *Amorphophallus hewittii*'s inflorescences. (A) Stingless bees in the pistillate zone. (B) Stingless bees on staminate zone. (C) Stingless bees. (D) Scavenger scarab beetle (Hybosoridae) (E-F) Scavenger scarab beetle on pistillate flowers. (G) Scavenger scarab beetles mating onto pistillate zone. (H) Fruit flies on pistillate flowers. (I) Green bottle flies (Calliphoridae). (J) Rove beetle (Staphylinidae) moving between pistillate flowers. (K) Unknown rove beetles on wet spadix. 72
- Figure 4.10 *Amorphophallus julaiihii* (A) Whole plants. (B) Immature infructescence. (C) Ripened infructescence. 73
- Figure 4.11 Inflorescence of *Amorphophallus julaiihii* at different stages. (A) Pistillate phase (left). (B) Staminate phase (left). (C-D) Rove beetles (Staphylinidae) at pistillate zone, some move on pistillate flowers. (E-F) Rove beetles collecting pollen on stamens. 74
- Figure 4.12 *Anadendrum* sp. nov. (A) Mature plant with inflorescences. (B) Infructescence with fruits ripening from tip. (C) *Chabria* beetles chewing on ripened fruits. 81
- Figure 4.13 Inflorescence of *Anadendrum* sp. nov. at various stages. (A) Pistillate anthesis (1<sup>st</sup> day). (B) Pistillate anthesis (2<sup>nd</sup> day) (C) Post- 82

pistillate anthesis (3<sup>rd</sup> day). (D) Staminate anthesis. (E) Thrips (Thripidae) and rove beetles onto the spadix. (F) Thrip consuming stigmatic secretion. (G) Rove beetle (Staphylinidae). (H) Thrip collecting pollen.

- Figure 4.14 *Bucephalandra muluensis* (A) Mature plant in natural habitat. (B) Immature infructescence. (C) Ripened infructescence, fruits split to reveal seeds. (D-E) Seeds germinated in lower spathe. 85
- Figure 4.15 Inflorescence of *Bucephalandra muluensis* at various stages (A-C) Spathe expands during pistillate anthesis, flies move in through tiny gap at tip of the spathe. (D) *Colocasiomyia* flies (Drosophilidae) active in the pistillate zone. (E-F) Spathe expanded and upper spathe abscised during staminate anthesis. (G) Flies on pistillate flowers. (H) Pollen released. (I-J) *Colocasiomyia* flies (Drosophilidae) collecting pollen in the staminate zone. 86
- Figure 4.16 *Homalomena ardua* (A) Mature plant with inflorescences. (B) Inflorescences and infructescences. 91
- Figure 4.17 Inflorescence of *Homalomena ardua* at various stages. (A) Pistillate anthesis. (B) Staminate anthesis. (C) *Chaloenus* beetles (Chrysomelidae) chewing on staminate zone before pollen release. (D) *Chaloenus* beetles moving on pistillate flowers, and consuming sterile interstices and interpistillar staminodes. (E) *Colocasiomyia* flies (Drosophilidae) moving on pistillate flowers. (F) *Parastasia* 92

beetles (Scarabaeidae). (G) *Chaloenus* beetles on pistillate zone (top view). (H) *Parastasia* beetles chewing sterile interstices. (I) *Parastasia* beetles consuming pollen. (J) *Chaloenus* beetles and *Colocasiomyia* flies collecting pollen.

- Figure 4.18 *Homalomena ovatifolia* (A) Mature plant with inflorescences, red leaf form behind green leaf form. 93
- Figure 4.19 Inflorescence of *Homalomena ovatifolia* at various stages (A-C) 94  
Pistillate anthesis, spathe gap opened. (B) *Colocasiomyia* flies (Drosophilidae) moving on pistillate flowers. (C) *Colocasiomyia* flies mating on the spadix. (D) Post-pistillate anthesis, spathe closed back. (E) Staminate anthesis with pollen released. (F) *Colocasiomyia* flies on the pistillate zone (zoom). (G-H) *Colocasiomyia* flies in inflorescence destroyed by *Chaloenus* beetles. (I) *Chaloenus* beetles (Chrysomelidae) consuming the staminate zone before the staminate anthesis.
- Figure 4.20 *Homalomena rostrata* (A) Mature plant with inflorescence buds. 95  
(B) Inflorescences and infructescences at different stages. (C) Infructescences bending downwards. (D) Ripened infructescences with red fruits.
- Figure 4.21 Inflorescence of *Homalomena rostrata* at various stages. (A, B & 96  
D) Pistillate anthesis, spathe opened, flies attracted. (C) *Parastasia* beetles (Scarabaeidae) on pistillate flowers, consuming

interpistillar staminodes. (E) Staminate anthesis, spathe closing back. (F) *Parastasia* beetles consuming pollen. (G) *Colocasiomyia* flies (Drosophilidae) moving on pistillate flowers, consuming stigmatic secretion. (H) *Colocasiomyia* flies collecting pollen.

- Figure 4.22 *Homalomena sanguineovaginata* sp. nov. (A) Mature plant with inflorescences. (B) Inflorescence at pistillate anthesis visited by insects. 97
- Figure 4.23 Inflorescence of *Homalomena sanguineovaginata* sp. nov. at various stages. (A-B) Pistillate anthesis, spathe opened, flies attracted. (C) Staminate anthesis, pollen released. (D) *Parastasia* beetle, *Chaloenus* beetle and *Colocasiomyia* flies at the level of the spathe constriction. (E) *Colocasiomyia* flies on pistillate flowers. (F) *Colocasiomyia* flies collecting pollen. 98
- Figure 4.24 *Lasia spinosa* (A-C) Inflorescence during pistillate anthesis. (D) Inflorescence during staminate anthesis, a gap opened at bottom of spathe after pollen released. (E) Flowers (Pistillate anthesis), stigmas orange colour. (F) Rove beetle (Aleocharinae, Staphylinidae). (G) Sap beetle (Nitidulidae) near the narrow gap. (H) Flowers releasing pollen (Staminate anthesis), stigmas deep orange colour. (I) Snout and bark beetle (Curculionidae). (J) Rove beetle (Aleocharinae, Staphylinidae) (K) Sap beetles collecting 105

pollen. (L) Developing infructescence. (M) Seedlings growing on the same spot.

Figure 4.25 Inflorescence of *Lasia spinosa* at various stages. (A) Rove beetles (Aleocharinae, Staphylinidae) moving on flowers during the pistillate anthesis. (B) Rove beetle and sap beetle. (C-D) Sap beetles (Nitidulidae). (E) Ambrosia beetles (Scolytinae). (F-G) Rove beetle collecting pollen. (H) Ambrosia beetle on Sap beetle. (I) Silvanid beetle (Silvanidae). (J) Plant bug (Miridae). (K) Sap beetle collecting pollen. (L) Plant bug collecting pollen. (M) Rove beetles and Ambrosia beetles collecting pollen. (N) Sap beetles and rove beetles collecting pollen. (O) Sap beetles, Ambrosia beetles and rove beetles collecting pollen.

Figure 4.26 *Pothos insignis* (A) Inflorescence and infructescence. (B) Inflorescence during staminate anthesis, flies (Drosophilidae) and *Chabria* beetles (Chrysomelidae) collecting pollen. (C) Nearly open inflorescence. (D) Seeds germinated on host plant. (E) Inflorescence during pistillate anthesis. (F) Inflorescence during 3<sup>rd</sup> day of staminate anthesis, stamens senesced after one day of pollen release. (G) Seeds germinated on ground. (H) Mature plant on host tree.

Figure 4.27 Inflorescence of *Pothos insignis* at various stages. (A-C) Pistillate anthesis. (A) Fruit flies (Drosophilidae) and Hover fly (Syrphidae) consuming stigmatic liquid. (B) Fruit flies, *Chabria* beetle and